

CORRESPONDENCES BETWEEN $K3$ SURFACES

FEDERICA GALLUZZI AND GIUSEPPE LOMBARDO

with an Appendix by Igor Dolgachev

ABSTRACT. In this paper we show that there is a correspondence between some $K3$ surfaces with non-isometric transcendental lattices constructed as a twist of the transcendental lattice of the Jacobian of a generic genus 2 curve. Moreover, we show the existence of a correspondence between a general $K3$ surface with $\rho = 17$ and a Kummer surface having transcendental lattices $\mathbf{Q}$ -Hodge isomorphic.

1. INTRODUCTION

In this paper we study the existence of correspondences between $K3$ surfaces $X(k, m, n)$, with $k, m, n \in \mathbf{N}$, Picard number 17 and transcendental lattices $T(k, m, n) \cong U(k) \oplus U(m) \oplus \langle -2n \rangle$. In the fundamental paper [Mu] Mukai showed that correspondences between $K3$ surfaces exist if the transcendental lattices are Hodge isometric over $\mathbf{Q}$. This construction holds if the Picard number of the surfaces is greater or equal to 11. Nikulin improved the result afterwards in [N2] obtaining the lower bound 5 for the Picard number.

The two authors are supported by EAGER - European Algebraic Geometry Research training network, Contract number HPRN-CT-2000-00099 and by Progetto di Ricerca Nazionale COFIN 2000 "Geometria delle Variet  Algebriche".

The aim of our work is to realize examples of $K3$ surfaces with transcendental lattice which are not Hodge isometric but such that a correspondence between them already exists. This in particular implies the existence of an algebraic cycle on the middle cohomology of the product of two surfaces arbitrarily chosen in the constructed family. More in details, in the first sections we recall some basic notions and results on lattices and correspondences. In Section 4 we consider a generic genus 2 curve and we show the existence of a correspondence between the Jacobian of the curve and a $K3$ surface with isomorphic transcendental lattice. Since this construction involves a second $K3$ surface whose transcendental lattice has quadratic form multiplied by 2, in Section 5 and 6 we generalize this first example. First, we construct $K3$ surfaces "twisting" each direct summand of the transcendental lattice of the Jacobian by natural numbers. Then we find correspondences between them using both Mukai's theorem and Shioda-Inose structures which translate the problem into a problem of looking for isogenies between abelian varieties.

In this way we prove in Theorem 6.3 that all the $K3$ surfaces $X(k, m, n)$ are in correspondence to each other.

Finally, in Theorem 6.5 we show the existence of a correspondence between a general $K3$ surface of Picard rank 17 and a Kummer surface of the same rank having transcendental lattices $\mathbf{Q}$ -Hodge isomorphic.

2. PRELIMINARY NOTIONS.

2.1. Definitions. A *lattice* is a free $\mathbf{Z}$ -module $\mathcal{L}$ of finite rank with a $\mathbf{Z}$ -valued symmetric bilinear form $b_{\mathcal{L}}(x, y)$. A lattice is called *even* if the quadratic form associated to the bilinear form has only even values, *odd* otherwise. A very useful invariant (under base change) of a lattice is its *discriminant* $d(\mathcal{L})$, defined as the determinant of the matrix of its bilinear form. Then, a lattice is called *non-degenerate* if the discriminant is non-zero and *unimodular* if the discriminant is ± 1 . If the lattice $\mathcal{L}$ is non-degenerate, the pair (s_+, s_-) , where $s_{\pm}$ denotes the multiplicity of the eigenvalue ± 1 for the quadratic form associated to $\mathcal{L} \otimes \mathbf{R}$, is called *signature* of $\mathcal{L}$. Finally, we call $s_+ + s_-$ the *rank* of $\mathcal{L}$ and $s_+ - s_-$ its *index*, moreover a lattice is *indefinite* if the associated quadratic form has both positive and negative values.

Given a lattice $\mathcal{L}$ we can construct the lattice $\mathcal{L}(m)$, that is the $\mathbf{Z}$ -module $\mathcal{L}$ with bilinear form $b_{\mathcal{L}(m)}(x, y) = mb_{\mathcal{L}}(x, y)$.

An *isometry* of lattices is an isomorphism preserving the bilinear form. Given a sublattice $\mathcal{L} \hookrightarrow \mathcal{L}'$, the embedding is *primitive* if $\frac{\mathcal{L}'}{\mathcal{L}}$ is free.

2.2. Examples.

- i) The lattice $\langle n \rangle$ is a free $\mathbf{Z}$ -module of rank one, $\mathbf{Z}\langle e \rangle$, with bilinear form

$$b(e, e) = n.$$

- ii) The *hyperbolic lattice* is the even, unimodular, indefinite lattice with $\mathbf{Z}$ -module

$$\mathbf{Z}\langle e_1, e_2 \rangle \text{ and bilinear associated form of matrix } \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

- iii) The lattice E_8 has $\mathbf{Z}^8$ as $\mathbf{Z}$ -module and the matrix of the bilinear form is the Cartan matrix of the root system of E_8 . It is an even, unimodular and positive definite lattice.

2.3. K3 and tori lattices. If X is a K3 surface, one can show that $H^2(X, \mathbf{Z})$ is free of rank 22 and that there is an isometry $H^2(X, \mathbf{Z}) \cong U^3 \oplus (E_8(-1))^2$. From now on we denote with Λ this K3-lattice. For X a complex torus, one has $H^2(X, \mathbf{Z}) \cong U^3$.

2.4. Hodge structures. Let X be an abelian or K3 surface. If we consider the Hodge decomposition of $H^2(X, \mathbf{C}) = H^{2,0}(X) \oplus H^{1,1}(X) \oplus H^{0,2}(X)$, inside $H^2(X, \mathbf{Z})$ one has two sublattices, the *Néron-Severi lattice*

$$NS(X) := H^2(X, \mathbf{Z}) \cap H^{1,1}(X)$$

and the orthogonal complement of $NS(X)$, the *transcendental lattice* T_X which has a natural Hodge structure induced by the one of $H^2(X, \mathbf{Z})$. The *Picard number* of X , denoted by $\rho(X)$, is the rank of $NS(X)$.

A *Hodge isometry* between the transcendental lattices of two K3 (or abelian) surfaces is an isometry preserving the Hodge decomposition.

2.5. Fourier-Mukai partners. Let X, Y be both abelian (or K3) surfaces, they are called *Fourier-Mukai partners* if their derived categories of bounded complexes of coherent sheaves are equivalent. Mukai and Orlov have showed that

Theorem 2.1. [Mu, Or] *X and Y as above are Fourier-Mukai partners if and only if their transcendental lattices are Hodge-isometric.*

3. KNOWN RESULTS.

In this section we recall some fundamental results which will play a key role in the next paragraphs. First of all, since for a $K3$ or an abelian surface the period map is surjective, it is possible to prove the following

Theorem 3.1. [Mor, Corollary 1.9] *Suppose $T \hookrightarrow U^3$ (resp. $T \hookrightarrow \Lambda$) is a primitive sublattice of signature $(2, 4 - \rho)$ (resp. $(2, 20 - \rho)$). Then there exists an abelian surface (resp. algebraic $K3$ surface) X and an isometry $T_X \xrightarrow{\sim} T$.*

Definition 3.2. A $K3$ surface X admits a *Shioda-Inose structure* if there is an involution ι on X such that $\iota^*(\omega) = \omega$ for every $\omega \in H^{2,0}(X)$ (ι is called a *Nikulin involution*) and with rational quotient map $\pi : X \dashrightarrow Y$ where Y is a Kummer surface. The map π_* induces a Hodge isometry $T_X(2) \cong T_Y$. This gives a diagram

$$\begin{array}{ccc} X & & Z \\ & \searrow \quad \swarrow & \\ & Y & \end{array}$$

of rational maps of degree 2 where Z is a complex torus and Y is the Kummer surface of Z .

One can detect the existence of a Shioda-Inose structure on a $K3$ surface analyzing the transcendental lattice of the surface.

Theorem 3.3. [Mor, Theorem 6.3] *Let X be an algebraic $K3$ surface. X admits a Shioda-Inose structure if and only if there is a primitive embedding $T_X \hookrightarrow U^3$.*

Remark 3.4. Obviously, the Shioda-Inose structure realizes a correspondence between the $K3$ surface X and the abelian variety Z ; this correspondence will play a fundamental role in the following paragraphs.

4. STARTING PROBLEM.

Let C be a generic genus 2 curve (i.e. such that its Jacobian surface has $\rho(JC) = 1$); JC is a principally polarized abelian variety and, if E is the principal polarization, so we have $E^2 = 2$ then $T := T_{JC} = U^2 \oplus \langle -2 \rangle$. Since we have obvious primitive embeddings of T in Λ , from Theorem 3.1 there exists an algebraic $K3$ surface X_1 such that $T_{X_1} \cong T_{JC}$. Moreover, by [Mu, Proposition 6.2], the number of Fourier-Mukai partners of X_1 is one since $\rho(X_1) = 17$, so such a X_1 is unique (up to isomorphisms). We analyze now the relations between the Jacobian surface and this $K3$ surface. First, we observe that we can construct an embedding $T_{X_1} \hookrightarrow U^3$ in the following way; we send the first two copies of $U \subset T_{X_1}$ to the corresponding ones of U^3 and the element of square -2 to $e_1^3 - e_2^3$ where $\{e_j^i\}_{j=1,2}^{i=1,2,3}$, is a basis of U^3 .

Then, from Theorem 3.3 X_1 admits a Shioda-Inose structure

$$\begin{array}{ccc} X_1 & & Z \\ & \searrow \quad \swarrow & \\ & X_2 & \end{array}$$

and one has $T_Z \cong T_{X_1}$, thus $T_Z \cong T_{JC}$. Recently, Hosono, Lian, Oguiso and Yau have proved the following

Theorem 4.1. [HLOY2, Main Theorem, 1)] *Let A and B abelian surfaces, they are Fourier-Mukai partners if and only if $Kum(A) \cong Kum(B)$.*

It follows that $Kum(Z) \cong Kum(JC)$ since Z and JC are Fourier-Mukai partners. In this way we obtain a correspondence between X_1 , X_2 and JC

$$\begin{array}{ccccc}
 JC & & Z & & X_1 \\
 & \searrow & & \searrow & \swarrow \\
 & Kum(JC) & \cong & X_2 = Kum(Z) &
 \end{array}$$

Remark 4.2. We observe that the Hodge isometry between T_{JC} and T_Z can be extended to a Hodge isometry between the second cohomology groups since JC and Z are principally polarized abelian surfaces of rank one. From [Sh], we obtain that $Z \cong JC$ or $Z \cong (JC)^\vee$.

5. EMBEDDINGS OF TWISTED LATTICES.

We want to generalize the situation of the previous paragraph. We consider the lattice $T(k, m, n) := U(k) \oplus U(m) \oplus \langle -2n \rangle$, with $k, m, n \in \mathbf{N}$, obtained by twisting the summands of T , equipped with the Hodge structure induced by T itself. This is an indefinite even lattice of signature $(2, 3)$. The aim is to prove the existence of correspondences between $K3$ surfaces having transcendental lattice Hodge-isometric to $T(k, m, n)$. In order to construct such surfaces, we start proving the following

Lemma 5.1. *The lattice $T(k, m, n)$ is a primitive sublattice of the $K3$ -lattice Λ .*

Proof. First we observe that a lattice $\langle -2t \rangle$, $t \in \mathbf{N}$, can be primitively embedded in $E_8(-1)$ from [N1, Theorem 1.12.4]. We denote by $\theta_t \in E_8(-1)$ its generator.

In order to construct the required primitive embedding of $T(k, m, n)$ in Λ we consider a θ_k contained in the first copy of $E_8(-1)$ and a θ_n in the second one.

Then, if $\{\delta_1, \dots, \delta_5\}$ is the standard basis of $T(k, m, n)$, we can obtain the desired embedding in the following way:

$$T(k, m, n) \hookrightarrow U^3 \oplus E_8(-1)^2$$

$$\delta_1 \mapsto (1, k) + (0, 0) + (0, 0) + \theta_k$$

$$\delta_2 \mapsto (1, 0) + (0, 0) + (0, 0)$$

$$\delta_3 \mapsto (0, 0) + (1, 0) + (0, m)$$

$$\delta_4 \mapsto (0, 0) + (0, 0) + (1, 0)$$

$$\delta_5 \mapsto \theta_n.$$

□

Remark 5.2. This Lemma, together with Theorem 3.1, implies the existence, for any $k, m, n \in \mathbf{N}$, of a $K3$ surface $X(k, m, n)$ with a Hodge isometry between the transcendental lattices $T_{X(k, m, n)}$ and $T(k, m, n)$. Such a surface is unique (up to isomorphisms) by [Mu, Proposition 6.2].

In order to generalize the example constructed in the previous paragraph, we want to produce correspondences among these $K3$ surfaces. We observe that, with these

notations, the surfaces of the previous section can be rewritten as $X_1 = X(1, 1, 1)$ and $X_2 = X(2, 2, 2)$.

The problem of finding correspondences among $K3$ surfaces was investigated by Mukai in [Mu]. His idea was to construct such correspondences starting from Hodge isometries (over $\mathbf{Q}$) between the transcendental lattices of the surfaces. This method works if the Picard number of the surfaces is sufficiently large. Mukai's result is the following

Theorem 5.3. [Mu, Cor.1.10.] *Let X, Y be $K3$ surfaces with $\rho(X), \rho(Y) \geq 11$. If $\varphi : T_X \otimes \mathbf{Q} \rightarrow T_Y \otimes \mathbf{Q}$ is a Hodge isometry, then φ is induced by an algebraic cycle.*

Our $K3$ surfaces, unfortunately, do not satisfy the condition of Mukai's theorem, since an isometry between $T(k, m, n)$ and $T(k', m', n')$ doesn't exist, even over $\mathbf{Q}$. However, in the same article Mukai proved the Oda's conjecture, as modified by Morrison in [Mor], which realizes a correspondence between a $K3$ surface X and an abelian surface provided that the $\mathbf{Q}$ -transcendental lattice $T_X \otimes \mathbf{Q}$ admits an embedding in $U^3 \otimes \mathbf{Q}$. In order to construct the correspondences among the $X(k, m, n)$, we follow a similar approach, translating the problem in another one involving some (abelian) surfaces which are in correspondence with the given ones.

6. ABELIAN SURFACES AND CORRESPONDENCES.

Motivated by the work of Morrison and Mukai, we start with the following

Lemma 6.1. *There is an embedding of $\mathbf{Q}$ -lattices $\phi : T(k, m, n) \otimes \mathbf{Q} \hookrightarrow U^3 \otimes \mathbf{Q}$.*

Proof. Let $\{e_1^i, e_2^i\}$, $i = 1, 2, 3$ be the basis of the i -th copy of the hyperbolic lattice U and let $\{a_1, b_1, a_2, b_2, c\}$ be the basis of $T(k, m, n)$. We define ϕ in the following way

$$a_1 \longmapsto e_1^1 \otimes 1$$

$$b_1 \longmapsto e_2^1 \otimes k$$

$$a_2 \longmapsto e_1^2 \otimes 1$$

$$b_2 \longmapsto e_2^2 \otimes m$$

$$c \longmapsto e_1^3 \otimes 1 + e_2^3 \otimes (-n).$$

□

The existence of such an embedding allows us to prove the following

Theorem 6.2. *For any $k, m, n \in \mathbf{N}$ there exist abelian surfaces A_n and correspondences*

$$\begin{array}{ccc} Z(k, m, n) & \longrightarrow & X(k, m, n) \\ \downarrow & & \\ A_n & & \end{array}$$

Proof. Let us consider the embedding $\phi : T(k, m, n) \otimes \mathbf{Q} \hookrightarrow U^3 \otimes \mathbf{Q}$ of Lemma 6.1 and the lattice $T_n := \phi(T(k, m, n) \otimes \mathbf{Q}) \cap U^3 = U^2 \oplus \langle -2n \rangle$ with the Hodge structure

induced by ϕ . Since T_n is a primitive sublattice of Λ , there exists a $K3$ surface Y_n and a Hodge isometry $T_{Y_n} \cong T_n$. Consider now the basis of $T_{X(k,m,n)}$ given in Lemma 6.1: the multiplication by $\frac{1}{k}$ on the sublattice $\mathbf{Z}\langle b_1 \rangle$ and by $\frac{1}{m}$ on $\mathbf{Z}\langle b_2 \rangle$ induces a Hodge isometry $T_{X(k,m,n)} \otimes \mathbf{Q} \cong T_{Y_n} \otimes \mathbf{Q}$ and Theorem 5.3 gives a correspondence $Z'(k, m, n)$ between $X(k, m, n)$ and Y_n .

We can define also a primitive embedding $T_n \cong T_{Y_n} \hookrightarrow U^3$ sending $U^2 \subset T_n$ to the first two copies of U in U^3 via the identity and sending the element of square $-2n$ to $e_1^3 - ne_2^3$. By Theorem 3.3 the existence of such an embedding is equivalent to the existence of a Shioda-Inose structure on the $K3$ surface Y_n . So there exist an abelian surface A_n , a Hodge isometry $T_{A_n} \cong T_n$ and a correspondence Z''_n between Y_n and A_n . The composition of the two correspondences

$$\begin{array}{ccccc}
 & Z'(k, m, n) & & Z''_n & \\
 & \swarrow & \searrow & \swarrow & \searrow \\
 X(k, m, n) & & Y_n & & A_n
 \end{array}$$

gives the desired one. □

Now, we are able to prove the following

Theorem 6.3. *Let $X(k, m, n)$, $k, m, n \in \mathbf{N}$, be a $K3$ surface with transcendental lattice Hodge-isometric to $T(k, m, n)$. For any $k', m', n' \in \mathbf{N}$ there exist correspondences*

$$\begin{array}{ccc}
Z_{k,m,n}^{k',m',n'} & \longrightarrow & X(k,m,n) \\
\downarrow & & \\
X(k',m',n') & &
\end{array}$$

and every $X(k,m,n)$ has a correspondence with the Jacobian surface JC of Section 4.

Proof. Let Z be a principally polarized abelian surface with $\rho(Z) = 1$ and let $\{\lambda_1, \lambda_2, \lambda_3, \lambda_4\}$ be a symplectic basis for the lattice of Z . Let now, for any n , $B_n := \mathbf{C}^2/\Lambda_n$ be the complex torus with lattice $\Lambda_n = \mathbf{Z}\langle\lambda_1, n\lambda_2, \lambda_3, \lambda_4\rangle$; the polarization E defines on each B_n a polarization of type $(1, n)$; moreover the abelian surfaces B_n are all obviously isogenous to each other.

On the other hand, one has that $T_{B_n} \cong U^2 \oplus \langle -2n \rangle$, so every B_n is a Fourier-Mukai partner of the surface A_n constructed in Theorem 6.2. This means, by Theorem 4.1, that $Kum(A_n) \cong Kum(B_n)$, thus there is a correspondence between A_n and B_n . In this way we have constructed a correspondence between A_n and A_m , for any n, m :

$$\begin{array}{ccccccc}
A_n & & B_n & \xrightarrow{\sim} & B_m & & A_m \\
\downarrow & & \downarrow & & \downarrow & & \downarrow \\
Kum(A_n) & \cong & Kum(B_n) & & Kum(B_m) & \cong & Kum(A_m)
\end{array}$$

But Theorem 6.2 gives a correspondence between $X(k,m,n)$ and A_n for any n , so the desired $Z_{k,m,n}^{k',m',n'}$ is the composition of these correspondences. Moreover, there is a correspondence between $X(1,1,1)$ and JC from Section 4, thus the statement follows.

□

Remark 6.4. We observe that the existence of the correspondence is independent from the chosen Hodge structure on T . However, it is necessary not to change the structure when we "twist" the starting lattice T by (k, m, n) . This allows us to obtain, for any chosen Hodge structure H on T , a family of $K3$ surfaces $X_H(k, m, n)$ in correspondence to each other.

Morrison, in [Mor, Cor 4.4] showed that a $K3$ surface K of Picard rank 17 is a Kummer surface if and only if there is an even lattice T' with $T_K \cong U(2)^2 \oplus T'(2)$. Then, this allows us to prove the following theorem, which generalizes Theorem 6.3

Theorem 6.5. *Let X be a general $K3$ surface with $\rho(X) = 17$ and $T_X \otimes \mathbf{Q}$ isomorphic as a Hodge structure to $T_K \otimes \mathbf{Q}$, where K is a Kummer surface. Then, there is a correspondence between X and K .*

Proof. Let $\psi_K = U^2(2) \oplus \langle -4n \rangle$ be the polarization on T_K , we show that there is a $\mathbf{Q}$ -basis of $T_X \otimes \mathbf{Q}$ in which the polarization is $\psi_X = a\psi_K$ with $a \in \mathbf{Q}$. Since $\psi_X \in \text{Sym}^2(T_X \otimes \mathbf{Q})^{MT(T_X \otimes \mathbf{Q})}$ (where MT denotes the Mumford-Tate group) it suffices to show that this space has dimension one. This is a consequence of Schur's Lemma, since $MT(T_x \otimes \mathbf{Q})(\mathbf{C}) = SO(5)(\mathbf{C})$ and the action on the Hodge structure is irreducible. So, we have the Hodge $\mathbf{Q}$ -isometries $T_X \otimes \mathbf{Q} \cong a(U^2(2) \oplus \langle -4n \rangle) \cong U^2 \oplus \langle -4na_1a_2 \rangle$, where $a = \frac{a_1}{a_2}$. From the surjectivity of the period map, the last one is a transcendental lattice of a $K3$ surface $X(1, 1, -4na_1a_2)$, which is in correspondence with X from Mukai's work. The statement now follows from Theorem 6.3. $\square$


ACKNOWLEDGMENTS

We would like to thank Prof. Bert van Geemen for having introduced us to this problem and for the useful discussions and valuable suggestions. We also thanks Prof. Igor Dolgachev for his interest in this work.

APPENDIX A. A GEOMETRIC CORRESPONDENCE BETWEEN X_1 AND A KUMMER SURFACE

We use the following result of V. Nikulin [N3].

Theorem A.1. *Let X be a K3-surface with $T_X \cong U \oplus \langle -2 \rangle$. Then $\text{Pic}(X)$ has only finitely many smooth rational curves which form the following graph.*



The automorphism group of X is generated by two commuting involutions σ and τ . The involution σ has 8 isolated fixed points. The set of fixed points of the involution τ is the union of 8 smooth rational curves and a smooth curve of genus 2.

It follows from observing the graph that X admits an elliptic fibration $|F|$ with two singular fibres

$$F_1 = 3R_0 + 2R_1 + 4R_2 + 6R_3 + 5R_4 + 4R_5 + 3R_6 + 2R_7 + R_8$$

and

$$F_2 = 2N_0 + N_1 + 2N_2 + 3N_3 + 4N_4 + 3N_5 + 2N_6 + N_7$$

of type $\tilde{E}_8$ and $\tilde{E}_7$. It is also has a section S . The fixed locus of τ consists of smooth rational curves $R_1, R_3, R_5, R_7, N_2, N_4, N_6, S$ and a genus 2 curve W which intersects R_0, N_0, N_7 with multiplicity 1. Let $p = W \cap R_0, q = W \cap N_0, a = W \cap N_7$. We have $3p \sim 2q + a$ and the fibration defines a g_3^1 on W spanned by the divisors $3p$ and $2q + a$.

We also observe that X contains another elliptic fibration $|F'|$ with two reducible fibres

$$F'_1 = 3N_0 + R_8 + 2S + 3N_1 + 4N_2 + 5N_3 + 6N_4 + 4N_5 + 2N_6$$

and

$$F'_2 = 2R_0 + A + 2R_1 + 3R_2 + 4R_3 + 3R_4 + 2R_5 + R_6$$

of type $\tilde{E}_8$ and $\tilde{E}_8$. The curve R_7 is a section. The involution σ switches the two fibres and induces the hyperelliptic involution on W . Its set of fixed points are 2 points on the curve R_8 and 6 points on W . Also note that σ maps the fibration $|F|$ to the fibration $|F'|$. It is easy to see that $q = \sigma(p)$. This gives $3p \sim 2K_W - 2p + a$, hence

$$|5p| = |2K_W + a|.$$

Theorem A.2. *The linear system $|F' + F|$ defines a map $f : X \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$ of degree 2. Its branch locus is a curve of bidegree $(4, 4)$ which is equal to the union of a curve B of bidegree $(3, 3)$ and two rulings E_1, E_2 . The curve B has 2 cusps q_1, q_2 . The cuspidal tangent at q_i is equal to E_i . The ramification curve of f is equal to W . The*

automorphism τ of X is the deck transformation of f , the quotient $X/(\tau)$ admits a birational morphism to $\mathbb{P}^1 \times \mathbb{P}^1$ which resolves the singularities of the branch curve. The automorphism σ is induced by the automorphism $\bar{\sigma}$ of $\mathbb{P}^1 \times \mathbb{P}^1$ which switches the factors. It leaves the curve B invariant and switches E_1 and E_2 .

Proof. We have $(F + F')^2 = 4$ and the restriction of $|F' + F|$ to a nonsingular fibre of each fibration is a degree 2 map. This easily implies that the linear system defines a degree 2 map f to a nonsingular quadric in $\mathbb{P}^3$ and the pre-images of the rulings are the fibrations $|F|$ and $|F'|$. The map f blows down the curves $R_0, R_1, R_2, R_3, R_4, R_6, N_0, N_1, N_2, N_3, N_4, N_5, N_6$. Its restriction to W is a birational map defined by a 3-dimensional linear subsystem of $|3p + 3q| = |3K_W|$. The rest of the assertions are easy to verify. $\square$

Consider the automorphism $\bar{\sigma}$ of $\mathbb{P}^1 \times \mathbb{P}^1$ and let

$$\pi : \mathbb{P}^1 \times \mathbb{P}^1 \rightarrow \mathbb{P}^1 \times \mathbb{P}^1 / (\bar{\sigma}) \cong \mathbb{P}^1$$

be the natural projection to the orbit space. Its locus of fixed points is the diagonal Δ . The image of Δ on $\mathbb{P}^2$ is a conic Q . The image of W is a cuspidal cubic G . The images of E_1 and E_2 is the cuspidal tangent T of G . It is also a tangent of the conic Q . The curves Q and G intersect at 6 points, the ramification points of the hyperelliptic involution of W .

Theorem A.3. *Let $\bar{f} : Y \rightarrow \bar{Y} \rightarrow \mathbb{P}^2$ be a minimal resolution of the double cover $\bar{Y}$ of $\mathbb{P}^2$ branched along the union of the curves G, Q and T . Then Y is a Kummer surface birationally isomorphic to the quotient of X by σ .*

Proof. We have a commutative diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & \mathbb{P}^1 \times \mathbb{P}^1 \\ \downarrow / \sigma & & \downarrow / \bar{\sigma} \\ Y & \xrightarrow{\bar{f}} & \mathbb{P}^2 \end{array}$$

It follows from [Na] that Y is a Kummer surface of $\text{Jac}(C)$ for some curve C (not isomorphic to W). □

Remark A.4. *One can reverse the construction of Theorem A.2. Starting from a genus 2 curve W together with a point $p \in W$ such that $|3K_W - p| \neq \emptyset$ and p is not a Weierstrass point, we construct a 2-cuspidal model $\bar{W}$ in $\mathbb{P}^1 \times \mathbb{P}^1$ as in the theorem. Then taking the double cover we get a K3 surface X with Picard lattice containing $U \oplus E_8 \oplus E_7$ and $U \oplus \langle -2 \rangle \subset T_X$. Replacing p by the conjugate point q under the hyperelliptic involution, we get the same surface X . As was explained to me by J. Harris, the number of pairs (p, q) as above is equal to 16. Thus we obtain that the moduli space of K3 surfaces marked with the lattice $U \oplus E_8 \oplus E_7$ is isomorphic to a $(16 : 1)$ -cover $\mathcal{M}'_2$ of the moduli space $\mathcal{M}_2$ of genus 2 curves, and on the other hand, via periods it is isomorphic to the moduli space $\mathcal{A}_2$ of principally polarized abelian surfaces. This defines a birational isomorphism between $\mathcal{M}'_2$ and $\mathcal{M}_2$.*

REFERENCES

- [B-M] T. Bridgeland, A. Maciocia, *Complex surfaces with equivalent derived categories*, Math.Z. **236** 4 (2001) 677-697
- [HLOY1] S. Hosono, B. H. Lian, K. Oguiso, S. Yau, *Counting Fourier-Mukai partners and applications*, alg-geom 0202014
- [HLOY2] S. Hosono, B. H. Lian, K. Oguiso, S. Yau, *Kummer structures on a K3 surface - An old question of T. Shioda*, alg-geom 0202082
- [L-B] C. Birkenhake, H. Lange, *Complex Abelian Varieties*. G.M.W. **302** Springer-Verlag, Berlin (1992)
- [Mor] D.R. Morrison, *On K3 surfaces with large Picard number*, Invent. Math. **75** 1 (1984) 105-121.
- [Mu] S. Mukai, *On the moduli space of bundles on K3 surfaces I*, in Vector Bundles on Algebraic varieties, M.F. Atiyah et al., Oxford University Press (1987), 341-413
- [Na] I. Naruki, *On metamorphosis of Kummer surfaces*, Hokkaido Math. Journal, **20**, (1991), 407-415.
- [N1] V. Nikulin *Integral symmetric bilinear forms and some of their applications*, Izv. Acad. Nauk. SSSR **43** (1979) 111-117 ; Math. USSR Izvestiya, **14** (1980) 1 103-167
- [N2] V. Nikulin *On correspondences between K3 surfaces*, Math. USSR Izvestiya, **30** (1988) 2 375-383
- [N3] V. Nikulin, *Quotient-groups of groups of automorphisms of hyperbolic forms by subgroups generated by 2-reflections. Algebro-geometric applications* (Russian) Current problems in mathematics, Vol. 18, pp. 3-114, Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Informatsii, Moscow, 1981 (Engl. Transl.: Journal of Soviet Math., **22**, (1983), 1401-1476).
- [Or] D.O. Orlov, *Equivalences of derived categories and K3 surfaces*, J. Math. Sci. (New York), **84** 5, (1997) 1361-1381, alg-geom 9606006

- [Sh] T. Shioda, *The period map of abelian surfaces*, J. Faculty Sci. Univ. Tokyo, Sect. 1A **25** 1 (1979) 47-59
- [Sh-I] T. Shioda, H. Inose, *On singular $K3$ surfaces*, in Complex analysis and algebraic geometry: papers dedicated to K.Kodaira. Iwanami Shoten and Cambridge University Press, (1977) 119-136

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI TORINO VIA CARLO ALBERTO 10 10123
TORINO, ITALY

E-mail address: federica.galluzzi@unito.it

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI TORINO VIA CARLO ALBERTO 10 10123
TORINO, ITALY

E-mail address: lombardo@dm.unito.it

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109, USA

E-mail address: idolga@umich.edu